

Name of the Topic: Hyperbolic function (Trigonometry)

## Expansion of Sinh x and Cosh x.

We have,  $\sinh x = i \sin ix$

$$= i \left[ ix - \frac{(ix)^3}{3!} + \frac{(ix)^5}{5!} - \dots \right]$$

$$= -i^2 \left[ x - \frac{i^2 x^3}{3!} + \frac{i^4 x^5}{5!} - \dots \right]$$

$$= x + \frac{x^3}{3!} + \frac{x^5}{5!} + \dots$$

Similarly,

$$\cosh x = \cos ix$$

$$= \left[ 1 - \frac{(ix)^2}{2!} + \frac{(ix)^4}{4!} - \dots \right]$$

$$= 1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \dots$$

## Periods of Hyperbolic func<sup>n</sup>

we have,

$$\sinh(\theta + 2n\pi i)$$

$$= -i \operatorname{Sinh}'(\theta + 2n\pi i)$$

$$= -i \operatorname{Sin}(i\theta - 2n\pi)$$

$$= +i \operatorname{Sin}(2n\pi - i\theta)$$

$$= -i \operatorname{Sin} i\theta = \sinh \theta.$$

Similarly,

$$\cosh(\theta + 2n\pi i)$$

$$= \cosh \theta \text{ and } \tanh(\theta + n\pi i)$$

$$= \tanh \theta.$$

Hence the hyperbolic sine and hyperbolic cosine are periodic, their periods being imaginary and equal to  $2n\pi i$ . The period of  $\tanh \theta$  is half of that of  $\sinh \theta$  and  $\cosh \theta$ .

Q. 2 Separate the following expression into their real and imaginary parts.

(i)  $\sin(x+iy)$

(ii)  $\cos(x+iy)$

(iii)  $\tan(x+iy)$

Co/3

$$\textcircled{1} \sin(x+iy) = \sin x \cosh y + i \cos x \sinh y$$

$$\textcircled{2} \cos(x+iy) = \cos x \cosh y - i \sin x \sinh y$$

$$\textcircled{3} \tan(x+iy) = \frac{\sin(x+iy)}{\cos(x+iy)}$$

$$\textcircled{4} \sin(2x) + \sin(2iy) = \sin 2x + i \sinh 2y$$

$$\textcircled{5} \cos(2x) + \cos(2iy) = \cos 2x + \cosh 2y$$

$$\textcircled{6} \sin^2 2x + \sin^2 2iy = \sin^2 2x + i \sinh 2y$$

$$\textcircled{7} \cos^2 2x + \cos^2 2iy = \cos^2 2x + \cosh 2y$$

Dr. P. Suresh Babu  
Asst. Prof. of Math.  
Dept. of Math.  
D.K. College,  
Dinduram.